

COMPUTABLY ENUMERABLE PARTIAL ORDERS

PETER A. CHOLAK, DAMIR D. DZHAFAROV, NOAH SCHWEBER,
AND RICHARD A. SHORE

ABSTRACT. We study the degree spectra and reverse-mathematical applications of computably enumerable and co-computably enumerable partial orders. We formulate versions of the chain/antichain principle and ascending/descending sequence principle for such orders, and show that the former is strictly stronger than the latter. We then show that every \emptyset' -computable structure (or even just of c.e. degree) has the same degree spectrum as some computably enumerable (co-c.e.) partial order, and hence that there is a c.e. (co-c.e.) partial order with spectrum equal to the set of nonzero degrees.

1. INTRODUCTION

A major theme of applied computability theory is the study of the algorithmic properties of countable structures and their presentations, and of the logical content of theorems concerning them. Partial orders, in particular, have been investigated extensively, most recently by Downey, Hirschfeldt, Lempp, and Solomon [3]; Hirschfeldt and Shore [6]; Jockusch, Kjos-Hanssen, Lempp, Lerman, and Solomon [7]; and Greenberg, Montalbán, and Slaman [4].

There are several approaches taken in such analyses. Most commonly, we restrict attention to computable orders and study the effectivity (or lack thereof) of particular combinatorial constructions or objects of interest. In computable model theory we might consider noncomputable orders, and inquire instead about which ones admit computable (isomorphic) copies, or more generally, in which Turing degrees copies can be found and how complicated the witnessing isomorphisms are. Finally, in reverse mathematics we formalize theorems pertaining to partial orders, and calibrate the strengths of these theorems according to which set-existence axioms are necessary to prove them. There is a fruitful interplay between these approaches, particularly the first and last, with results and insights from one often leading to results in the others. For a detailed account of this relationship, see for example [6, Section 1]. We refer the reader to Soare [12] for background in computability theory; to Ash and Knight [1] for computable model theory; and to Simpson [10] for reverse mathematics.

In this article, we look at several questions pertaining to computably enumerable (c.e.) and co-computably enumerable (co-c.e.) partial orders. Such orders, in which the relationship between a given pair of elements is not decidable but rather may only be revealed over time, arise naturally in different contexts in computability

The authors are grateful to Julia Knight and the anonymous referees for helpful comments. This research was partly conducted while Dzhafarov and Shore were visiting the Institute for Mathematical Sciences at the National University of Singapore in 2011, with funding from the John Templeton Foundation. Cholak was partially supported by NSF grant DMS-0800198, Dzhafarov by an NSF Postdoctoral Fellowship, and Shore by NSF grant DMS-0852811.

theory. For example, the inclusion order on a uniformly computable family of sets (possibly with repetition) is co-c.e., and in fact, in terms of computational complexity the two are interchangeable, as we show (Proposition 3.1). Previous work on c.e. partial orders was done by Case [2], who studied their extensibility to total orders, and Roy [9], who studied which c.e. binary relations have computable copies.

Our interest was motivated by generalizations of the chain/antichain principle (CAC) and the ascending/descending sequence principle (ADS); see Section 3 for definitions. In the context of reverse mathematics, where the orders one considers may be regarded as being computable, it is easy to see that CAC implies ADS over RCA_0 , but it is unknown whether the reverse implication holds. We show that when the principles are formulated for (formalizations of) c.e. and co-c.e. partial orders, the answer to the analogous question is no (Proposition 3.2 and Corollary 3.4).

We then look at the degree spectra of c.e. and co-c.e. partial orders on ω , the degree spectrum of an order being the class of degrees containing a copy of it. We show (Theorem 2.1) that even though there are c.e. partial orders with no co-c.e. copy, and co-c.e. ones with no c.e. copy, the degree spectra of the two classes of orders coincide (Corollary 4.4). Our main result is that the degree spectra of these partial orders are in fact universal for \emptyset' -computable structures: for every such structure (or indeed for any structure of c.e. degree) there is a c.e. (or co-c.e.) partial order with the same degree spectrum, and of the same degree if the structure has c.e. degree (Theorem 4.3). As a corollary, we obtain a new example of the Slaman-Wehner theorem ([11], [13]) that there is a structure with a copy in every nonzero degree. Specifically, it follows that such a structure can be found among the c.e. and co-c.e. partial orders in any nonzero c.e. degree (Corollary 4.6).

2. PRELIMINARIES


In this paper, we are interested in countably infinite structures and (unless otherwise noted) assume that their domains are ω . We also identify orders with the binary relations defining them. Our terminology and notation will for the most part be standard. Given a partial order \leq_P on ω and $A, B \subseteq \omega$, we say $A \leq_P B$ when $(\forall a \in A)(\forall b \in B)[a \leq_P b]$, and similarly for $A <_P B$. We write $A \not\leq_P B$ if $(\exists a \in A)(\exists b \in B)[a \not\leq_P b]$. When A or B is finite, we often replace it by its members, as in $A \not\leq_P b$ and $a \leq_P b_0, b_1$ in place of $A \not\leq_P \{b\}$ and $\{a\} \leq_P \{b_0, b_1\}$, respectively.

In referring to or building a c.e. partial order \leq_P on ω , we usually assume we begin with a computable partial order. We then enumerate additional pairs $\langle a, b \rangle$ into the order relation and speak of (computably) adding elements to (the graph of) \leq_P , or of setting $a \leq_P b$ for some $a, b \in \omega$. In the co-c.e. case, we again begin with some computable partial order, and then remove pairs, or set $a \not\leq_P b$. In both cases, of course, we must take care to build a transitive relation.

In this section we prove that c.e. and co-c.e. partial orders cannot be used interchangeably. The following theorem extends Theorem 2.5 of Roy [9] that there is a c.e. antisymmetric binary relation with no computable copy.

Theorem 2.1. *There exists a co-c.e. partial order on ω which is not isomorphic to any c.e. such order, and conversely.*

Proof. We partition ω computably into the following sets:


 FIGURE 1. The number i coded in \leq_P .

- $\{a, b, c, f, l\}$;
- $A = \{a_{i,k} : i \in \omega, k < i + 1\}$;
- $B = \{b_i : i \in \omega\}$;
- $C = \{c_{i,k} : i \in \omega, k < i\}$.

We shall use the elements of the first of these sets to identify the elements of the others, and the relations that hold between them, in a given copy of (ω, \leq_P) . For each $i \in \omega$, the $a_{i,k}$ for $k < i + 1$ will code the number i as a sequence of length $i + 1$. The elements f and l will identify the first and last element of this sequence, and the $c_{i,k}$ for $k < i$ will identify pairs of consecutive elements.

Let U be a fixed Σ_2^0 -complete subset of ω , and let R be a computable predicate so that $U = \{i : (\exists x)(\forall y)R(i, x, y)\}$. We use B to represent whether or not $U(i)$ holds, with $b_x \leq_P a_{i,0}, \dots, a_{i,i}$ if and only if we believe that $(\forall y)R(i, x, y)$ holds. Whenever we find some y such that $R(i, x, y)$ does not hold, we thus set $b_x \not\leq_P a_{i,0}, \dots, a_{i,i}$, so that in the end we have that $U(i)$ holds if and only if there is an $n \in B$ with $n \leq_P a_{i,0}, \dots, a_{i,i}$, namely $n = b_x$ for the least x such that $(\forall y)[R(i, x, y)]$.

Formally, we build \leq_P as follows. The initial setup is to remove elements from ω^2 so that no relations hold except for those needed for reflexivity and the following:

- $A <_P a, c$;
- $B <_P b$;
- $C <_P c$;
- $B <_P A$;

and for all $i \in \omega$,

- $a_{i,0} <_P f$ and $a_{i,i} \leq_P l$;
- $a_{i,k} <_P c_{i,k}$ and $a_{i,k+1} <_P c_{i,k}$ for all $k < i$;

See Figure 1 for an illustration.

We now proceed by stages, at each of which we remove at most finitely many elements from the graph of \leq_P . Associate to each $i \in \omega$ a natural number called its *witness*, initially declared to be 0. At stage s , consider each $i \leq s$ in turn, and suppose the witness of i is x . If $R(i, x, y)$ holds for all $y \leq s$, do nothing. Otherwise, set $b_x \not\leq_T a_{i,0}, \dots, a_{i,i}$, and redefine the witness of i to be $x + 1$.

The resulting order is clearly co-c.e. We also have that A is the set of elements strictly below a and not below b , B is the set of elements strictly below b , and C the set of elements strictly below c and not below a . Hence, the images of these sets in any copy of (ω, \leq_P) are computable in that copy, and if the order in that copy is c.e. then so are B and C .

Seeking a contradiction, suppose there exists a c.e. copy (ω, \leq_Q) of (ω, \leq_P) . Identify f, l, A, B , and C with their images in this copy. Given i , we search for elements $a_0, \dots, a_i \in A$ such that

- $a_0 \leq_Q f$ and $a_i \leq_Q l$;
- for each $k < i$, there is an element of C above both a_k and a_{k+1} .

This search is computable since \leq_Q and C are c.e. Furthermore, the search must succeed since the images in (ω, \leq_Q) of the elements $a_{i,0}, \dots, a_{i,i}$ from (ω, \leq_P) would do, and these are also the only possibilities. By construction, we thus have that $i \in U$ if and only if there is an $n \in B$ with $n \leq_Q a_0, \dots, a_i$. But this now yields a Σ_1^0 definition of U since B is c.e., contradicting the choice of U as a Σ_2^0 -complete set.

A similar argument yields a c.e. partial order with no co-c.e. copy. \square

While we are here particularly interested in partial orders, there are other natural binary relations one could consider. For example, partial orders can be viewed as directed graphs and we could obtain the preceding theorem for graphs by virtually the same argument. These classes of relations (and others) were studied in [5] alongside partial orders, and the same results were obtained for each. Similarly, we can recast all the results of Section 4 about spectra of c.e. and co-c.e. structures for any of these relational structures (see the comments at the end of Section 4).

3. GENERALIZATIONS OF ADS AND CAC

In this section, we investigate the reverse mathematical strength of several mild generalizations of the principles CAC and ADS, defined below. Recall that a *chain*, respectively, *antichain*, for a partial order \leq_P on an arbitrary set $S \subseteq \omega$ is a subset of S any two members of which are \leq_P -comparable, respectively, \leq_P -incomparable. An *ascending sequence*, respectively *descending sequence*, for \leq_P , is a subset of S on which \leq_P agrees with the natural order \leq_ω , respectively, the reverse natural order \geq_N .

The following principles are defined in the language of second-order arithmetic. By usual convention, we write \mathbb{N} in place of ω to refer to the set of natural numbers in an arbitrary model of RCA_0 .

Chain/antichain principle (CAC). *Every partial order on \mathbb{N} has either an infinite chain or an infinite antichain.*

Ascending/descending sequence principle (ADS). *Every linear order on \mathbb{N} has either an infinite ascending sequence or an infinite descending sequence.*

The reverse mathematical content of CAC and ADS was studied by Hirschfeldt and Shore [6], who established, among other results, that both principles are strictly weaker than Ramsey's theorem for pairs (RT_2^2) and incomparable with WKL_0 . It is easy to see that CAC implies ADS over RCA_0 . The question of the reverse implication, originally posed as Question 6.1 in [6], has been an open question in reverse mathematics for a number of years.¹ We show that the answer to the question is no when we consider more general types of orders.

To begin, we can easily formalize the notions of c.e. partial and linear orders in RCA_0 : a *c.e. partial order on \mathbb{N}* is a function, the elements of whose range, viewed as pairs, include $\langle i, i \rangle$ for all $i \in \mathbb{N}$, and additionally satisfy antisymmetry and transitivity; a *c.e. linear order on \mathbb{N}* is a c.e. partial order whose range includes $\langle i, j \rangle$ or $\langle j, i \rangle$ for all $i, j \in \mathbb{N}$. We define *co-c.e. partial orders* and *co-c.e. linear*

¹A negative answer was announced by Lerman, Solomon, and Towsner as this article went to press.

orders the same way, but referring, instead of to the range of a function, to the complement of the range. Chains and antichains, and ascending and descending sequences, are defined as before, provided i being below j is interpreted to mean that $\langle i, j \rangle$ is in the range (or complement of the range) of the relevant function.

With these definitions, we are able to formulate analogues of CAC and ADS for c.e. and co-c.e. orders:

CAC for c.e. (co-c.e.) partial orders on \mathbb{N} . *Every c.e. (co-c.e.) partial order on \mathbb{N} has either an infinite chain or an infinite antichain.*

ADS for c.e. (co-c.e.) linear orders on \mathbb{N} . *Every c.e. (co-c.e.) linear order on \mathbb{N} has either an infinite ascending sequence or an infinite descending sequence.*

Individual instances of these principles are thus no longer computable partial and linear orders, but rather c.e. and co-c.e. ones.

A different but related class of orders is that of inclusion orders on families of sets. For computable families, inclusion is co-c.e. But since a family of sets can in general contain repetitions, we do not necessarily obtain a co-c.e. partial order isomorphic to the inclusion order on a given computable family $\langle A_i : i \in \omega \rangle$ simply by setting $i \leq j$ if $A_i \subseteq A_j$. However, we do obviously obtain a preorder (a reflexive, transitive relation, not necessarily antisymmetric). The converse is also true:

Proposition 3.1. *Every co-c.e. preorder on ω is isomorphic to the inclusion order on a computable family of sets.*

Proof. Fix a co-c.e. preorder (ω, \leq_P) , along with a computable enumeration of its complement. Write $i \not\leq_{P,s} j$ and $i \leq_{P,s} j$ depending on whether the pair $\langle i, j \rangle$ has or has not been enumerated by stage s , and assume our enumeration has been speeded up, if necessary, to ensure that the relation $\leq_{P,s}$ is transitive on $\omega \upharpoonright s + 1$. We define $\langle A_i : i \in \omega \rangle$ by stages, defining the A_i for $i \leq s$ on a common initial segment of ω at stage s . The isomorphism will be given by $i \mapsto A_i$.

At stage s , add to A_s all elements of A_i for all $i \leq s$ with $i \leq_{P,s} s$. Then, consider any $i, j \leq s$ with $i \not\leq_{P,s} j$, and let n be the least number at which no set has yet been defined. For each $k \leq s$, add n to A_k if $i \leq_{P,s} k$, and to the complement of A_k otherwise.

The family $\langle A_i : i \in \omega \rangle$ is computable. It is also clear that if $i \not\leq_P j$ then $A_i \not\subseteq A_j$, as some number will be put into $A_i - A_j$ at a sufficiently large stage. Now suppose $i \leq_P j$, so that $i \leq_{P,s} j$ for all s . By our assumption about $\leq_{P,s}$, for any stage $s \geq \max\{i, j\}$ and any $k \leq s$, we will necessarily have $k \leq_{P,s} j$ whenever $k \leq_{P,s} i$. Thus, any number put into A_i at such a stage will also belong to A_j . Since numbers can only be put into A_i at a stage $s \geq i$, it thus follows that if $j < i$ then $A_i \subseteq A_j$. On the other hand, if $i < j$, then all elements of A_i are put into A_j at stage $s = j$, so we reach the same conclusion. \square

It is worth emphasizing that all of our preorders (c.e., co-c.e., or computable) below are defined on ω , and as such, equality of elements in the domain is still computable.

We can formalize c.e. and co-c.e. preorders on \mathbb{N} in RCA_0 just as we did partial orders, and then consider formulations of CAC and ADS for them as well. We can also look at versions for *actual* preorders, that is, those given as relations rather than ranges of functions. (We omit the statements, which are entirely analogous to the ones above.) While we could formulate versions of CAC and ADS directly for families of sets under inclusion, this would be unnecessary by the preceding

proposition (whose statement and proof can easily be formalized and carried out in RCA_0 .)

Each of the generalized principles can easily be seen to imply the corresponding original one. In the case of ADS, and in the case of CAC for preorders, we also obtain reversals:

Proposition 3.2. *Over RCA_0 , the following are equivalent:*

- (1) ADS;
- (2) ADS for c.e. linear orders on \mathbb{N} ;
- (3) ADS for co-c.e. linear orders on \mathbb{N} ;
- (4) ADS for c.e. linear preorders on \mathbb{N} ;
- (5) ADS for co-c.e. linear preorders on \mathbb{N} ;
- (6) ADS for linear preorders on \mathbb{N} .

In addition, CAC for preorders on \mathbb{N} is equivalent to CAC.

Proof. That (1) is implied by each of the other statements is clear. To see that (1) implies (2), note that, by linearity, the range of every c.e. linear order on \mathbb{N} exists. It is thus a linear order that we may apply ADS to, and any ascending or descending sequence for this order is such a sequence for the original c.e. order. The same argument shows that (1) implies (3).

Each of (4) and (5) implies (6), so it remains only to show that (1) implies (4) and (5). We prove the former, the proof of the latter being similar. So let a c.e. linear preorder be given. Formally, this is a function, but we denote it by \leq_P and write $i \leq_P j$ to mean that $\langle i, j \rangle$ is in the range. We write $i \sim_P j$ if $i \leq_P j$ and $j \leq_P i$. There are two cases to consider.

First, suppose there exist $i_0, \dots, i_{n-1} \in \omega$ such that $i_j \not\sim_P i_k$ for all distinct $j, k < n$, and for every i there is a $j < n$ such that $i \sim_P i_j$. Then for each $j < n$, the set $I_j = \{i \in \mathbb{N} : i \sim_P i_j\}$ exists by Δ_1^0 comprehension. By the infinitary pigeonhole principle (RT^1), which follows from ADS ([6, Proposition 4.5]), we may fix a $j < n$ such that I_j is infinite. Then this is an ascending (and also descending) sequence for \leq_P .

Now suppose no such i_0, \dots, i_{n-1} as above exist. Then for every finite set F there exists an $i > \max F$ such that $i \not\sim_P j$ for all $j \in F$, and, in fact, there is a function g , total by assumption, that assigns to each F the least such i . We now define by primitive recursion a function $f : \mathbb{N} \rightarrow \mathbb{N}$ such that $f(i) = g(f \upharpoonright i)$ for all $i \in \mathbb{N}$. Let R be the range of f , which exists because f is strictly increasing and is thus also infinite. For all $i, j \in R$ we have $i <_P j$ or $j <_P i$. If we write the elements of R as $r_0 < r_1 < \dots$, we can define a linear order $<_Q$ on \mathbb{N} by $i <_Q j$ if and only if $r_i <_P r_j$, which exists by Δ_1^0 comprehension. If S is any infinite ascending or descending sequence for (\mathbb{N}, \leq_Q) as given by ADS, then $\{r_i : i \in S\}$ is such a sequence for (\mathbb{N}, \leq_P) .

A similar argument establishes the equivalence of CAC for preorders on \mathbb{N} with CAC. (Of course, since a given preorder \leq_P is a relation in this case, the set of all $j \sim_P i$ for a given i exists even by Σ_0^0 comprehension.) \square

Essentially, the reason for the equivalences in the preceding proposition is that every c.e. or co-c.e. linear order is actually computable. By contrast, we have already seen in Theorem 2.1 that this is not the case for general partial orders. This also follows from the next theorem, as computable instances of CAC always have solutions that do not compute \emptyset' (see Diagram 1 in [6]).

Theorem 3.3.

- (1) *There exists a co-c.e. partial order on ω with no infinite antichains and with all chains computing \emptyset' .*
- (2) *There exists a c.e. partial order on ω with no infinite chains and with all infinite antichains computing \emptyset' .*

Proof. To prove (1), fix a computable enumeration $\langle \emptyset'_s : s \in \omega \rangle$ of \emptyset' , and let t_i be the least s such that $\emptyset'_s \upharpoonright i = \emptyset' \upharpoonright i$. We build a co-c.e. partial order \leq_P by stages. To begin, we make \leq_P agree with the natural order on ω .

At stage 0, we do nothing. At stage $s > 0$, we consider consecutive substages $i \leq s$. At substage i , if no number enters $\emptyset' \upharpoonright i$ at stage s , we do nothing and go either to substage $i + 1$ or to stage $s + 1$, depending on whether $i < s$ or $i = s$. Otherwise, for all j, k with $i \leq j < k \leq s$, we make $j \not\leq_P k$ and go to stage $s + 1$.

It is easily seen that \leq_P is a co-c.e., reflexive, antisymmetric relation, while its transitivity follows from the fact that $i \leq_P j$ implies $i \leq j$, and that if $i \not\leq_P k$ for some $k > i$ then necessarily $i \not\leq_P j$ for all j with $i < j \leq k$. So \leq_P is a partial order, and we note that it admits no infinite antichain since for every i , $i \leq_P j$ for all $j \geq t_i$.

Suppose now that $C = \{c_0 < c_1 < \dots\}$ is an infinite chain for \leq_P , so that $c_0 <_P c_1 <_P \dots$. In particular, since $i \leq c_i$, we must have $c_{i+1} \geq t_i$ for all i by construction. Thus for every i we have $\emptyset' \upharpoonright i = \emptyset'_{c_{i+1}} \upharpoonright i$, whence it follows that $\emptyset' \leq_T C$, as desired.

The argument for (2) is analogous. □

Formalizing the previous argument immediately yields the following (note that (4) and (5) are included since all partial orders are preorders):

Corollary 3.4. *Over RCA_0 , the following are equivalent:*

- (1) ACA_0 ;
- (2) CAC for c.e. partial orders on \mathbb{N} ;
- (3) CAC for co-c.e. partial orders on \mathbb{N} ;
- (4) CAC for c.e. preorders on \mathbb{N} ;
- (5) CAC for co-c.e. preorders on \mathbb{N} .

Together with Proposition 3.2, this provides a separation between the c.e. and co-c.e. forms of CAC on the one hand, and the c.e. and co-c.e. forms of ADS on the other.

4. DEGREE SPECTRA

The *degree spectrum* of a countable structure \mathcal{S} is the set of Turing degrees of copies of \mathcal{S} . The study of degree spectra, and in particular, of which classes of degrees can be realized as spectra, has been the subject of many investigations in computable model theory. A partial survey of previous results appears in Section 1 of [5].

Every structure \mathcal{S} we consider below will be assumed to be in a computable language, with computable signature. We also assume that \mathcal{S} is automorphically nontrivial (meaning there is no finite $F \subseteq |\mathcal{S}|$ such that each permutation of $|\mathcal{S}|$ fixing F is an automorphism), as trivial structures are structurally uninteresting. By Knight's theorem ([8], Theorem 4.1), the degree spectrum of any \mathcal{S} we consider

is therefore closed upwards, and thus also equal to the set of degrees of sets that compute a copy of \mathcal{S} .

In this section, we show that the degree spectra of c.e. and co-c.e. partial orders are universal for \emptyset' -computable structures. In contrast with Theorem 2.1, this also shows that the classes of degree spectra of c.e. and co-c.e. partial orders are the same.

We begin with a reduction of arbitrary structures to graphs (symmetric, irreflexive binary relations). As with orders, we identify graphs on ω with their relations.

Lemma 4.1. *For every structure \mathcal{S} with domain ω there exists a graph R on ω such that $R \equiv_T \mathcal{S}$ and R and \mathcal{S} have the same degree spectrum.*

Proof. This proof of this result is folklore, and has appeared in various forms in the published and unpublished literature. A complete proof can be found in [5, Appendix A]. \square

Lemma 4.2. *For every \emptyset' -computable graph R on ω there exists a c.e. (and a co-c.e.) partial order \leq_P on ω with the same degree spectrum. Furthermore, if R has c.e. degree then $\deg(\leq_P) = \deg(R)$.*

Proof. We begin by providing a uniform effective transformation taking any graph R to a partial order \leq_P with a uniformly effective inverse transformation taking any copy of this partial order to a copy of R . Thus R and \leq_P have the same degree spectrum. We partition ω computably into the following sets:

- $\{a, g, r_0, r_1\}$;
- $A = \{a_i : i \in \omega\}$
- $G = \{g_{i,j,k} : i < j, k \in \omega\}$.

Intuitively, the a_i represent the elements of the domain of a given copy of R and the $g_{i,j,k}$ represent guesses at whether or not R holds of the pair (a_i, a_j) in that copy. We code these guesses using r_0 and r_1 , with r_0 representing that R does not hold, and r_1 that it does. (The guessing will be tied to a computable approximation of R when $\deg(R)$ is c.e.) The numbers a and g help us recognize the pieces of the above partition in a given copy of \leq_P .

We can describe the isomorphism type of the desired partial order by saying that A is the set of elements below a and not below g , and G is the set of elements below a and g . For each $i < j$ and all k , $g_{i,j,k}$ is below a_i, a_j , and g . Additionally, all but one $g_{i,j,k}$ are below both r_0 and r_1 , and exactly one is below just one of r_0 and r_1 . The latter $g_{i,j,k}$ is below r_0 if $R(i, j)$ does not hold, and below r_1 otherwise. It is easy to see that we can build an order of this type computably in R , and that, conversely, any order of this type computes a copy of R (and indeed, that the transformation in either direction preserves degree). The point here is that, given a copy of this partial order, we start with the elements corresponding to a, g, r_0 and r_1 and then compute a copy of R with domain A by, for each $a_0, a_1 \in A$, searching for a $g \in G$ with $g \leq a_0, a_1$ such that $g \not\leq r_0$ or $g \not\leq r_1$. Exactly one of these must happen and we let there be an edge between a_0 and a_1 if and only if it is the first.

When R is \emptyset' -computable, we make \leq_P c.e. as follows. By Knight's theorem, as discussed above, we may assume that R has c.e. degree, since if not we can replace it by a copy of degree \emptyset' . Initially, we let \leq_P be the transitive closure of the following conditions:

- $A <_P a$;




FIGURE 2. A configuration in \leq_P . Since $g_{i,j,2}$ is below r_0 but not below r_1 , this codes that $R(i, j)$ does not hold.

- $G <_P g$;
- $g_{i,j,k} \leq_P a_i, a_j$ for all $i < j$ and k .

By the modulus lemma there is a computable function $f(i, j, s)$ with $\lim_s f(i, j, s) = 0$ if $R(i, j)$ does not hold and $\lim_s f(i, j, s) = 1$ if it does, and with modulus of convergence m of the same degree as R . At stage $s > 0$, for every $i < j \leq s$, we let k be the largest number less than or equal to s such that $f(i, j, k) \neq f(i, j, k-1)$, or 0 if there is no such number. We make $g_{i,j,k} \leq_P r_{f(i,j,k)}$ and $g_{i,j,l} \leq_P r_0, r_1$ for all $l \leq s$ not equal to k . See Figure 2 for an illustration.

It is easy to see from the limit properties of $f(i, j, s)$ that \leq_P is c.e. and that it has the isomorphism type described above. Thus, in particular, its degree spectrum is the same as that of R . Furthermore, for each $i < j$ it is precisely $g_{i,j,m(i,j)}$ which is below just one of r_0 and r_1 (indeed, it is below just $r_{f(i,j,m(i,j))} = r_{\lim_s f(i,j,s)}$, as desired). Thus, \leq_P is of the same degree as m , and so of the same degree as R .

To obtain instead a co-c.e. partial order, we modify the construction of \leq_P by initially setting $g_{i,j,k} \leq_P r_0$ and $g_{i,j,k} \leq_P r_1$ for all $i < j, k \in \omega$, and then removing one of these relations when our guess about $R(i, j)$ changes (including by default at $k = 0$) and both of them when it does not. In the end, there will be exactly one k such that $g_{i,j,k}$ is below r_0 or r_1 . It will be below just one of them and $R(i, j)$ will hold just in case it is below r_1 . This completes the proof. \square

Theorem 4.3. *For every \emptyset' -computable structure \mathcal{S} on ω there exists a c.e. (and a co-c.e.) partial order on ω with the same degree spectrum as \mathcal{S} . Furthermore, there exists such a partial order in every c.e. degree containing a copy of \mathcal{S} .*

Proof. By Lemma 4.1, for every \emptyset' -computable copy of \mathcal{S} on ω there is a graph R on ω of the same degree and with the same degree spectrum. By Lemma 4.2, there exists a c.e. (and a co-c.e.) partial order with the same degree spectrum as R , and if the degree of a given copy of \mathcal{S} , and hence of the corresponding one of R , is c.e., we may choose the order to be of that degree as well. \square

As an immediately corollary, we have:

Corollary 4.4. *Let \mathcal{S} be a structure with domain ω . The following are equivalent:*

- (1) \mathcal{S} has a copy of c.e. degree;
- (2) \mathcal{S} has a \emptyset' -computable copy;

- (3) *there is a c.e. (and a co-c.e.) partial order on ω with the same degree spectrum as \mathcal{S} .*

By Theorem 4.3, we obtain a wide array of classes of degrees realized as degree spectra of c.e. (or co-c.e.) partial orders, including, for example, the class of degrees $\geq \mathbf{0}'$. A particularly interesting class is that of the degrees strictly above $\mathbf{0}$. Recall the following well-known result:

Theorem 4.5 (Slaman [11], Wehner [13]). *There exists a countable structure whose degree spectrum consists precisely of the nonzero degrees.*

Since any two copies of the same structure have the same degree spectrum, it follows that an \mathcal{S} as above can be found in any nonzero degree. The following consequence of Theorem 4.3 is a considerably more effective version.

Corollary 4.6. *Every nonzero c.e. degree contains a c.e. (and a co-c.e.) partial order on ω whose degree spectrum consists precisely of the nonzero degrees.*

Proof. Fix a nonzero c.e. degree, and any structure in that degree satisfying Theorem 4.5. Then apply Theorem 4.3. \square

As partial orders are directed graphs, all the results of this section hold for directed graphs as well. As for (undirected) graphs, one could prove a general coding result taking c.e. or co-c.e. partial orders to c.e. or co-c.e. graphs as in [5, Appendix A]. There is a simpler route in our case, however. The crucial item here is Lemma 4.2. Note first that the comparability graph \tilde{R} of a partial order \leq_P , defined for x, y in the domain of \leq_P by $\tilde{R}(x, y) \iff x <_P y \vee y <_P x$, is always computable from \leq_P and is c.e. (co-c.e.) if \leq_P is. Next, given a graph R as in Lemma 4.2 and the corresponding partial order \leq_P computing R constructed there, it is clear that the associated comparability graph \tilde{R} has enough information to compute R , and hence also \leq_P . It is then a c.e. (co-c.e.) graph of the same degree, and with the same degree spectrum, as \leq_P . Thus all the results of this section also hold for graphs in place of partial orderings.

REFERENCES

- [1] C. J. Ash and J. Knight. *Computable structures and the hyperarithmetical hierarchy*, volume 144 of *Studies in Logic and the Foundations of Mathematics*. North-Holland Publishing Co., Amsterdam, 2000.
- [2] John Case. Sortability and extensibility of the graphs of recursively enumerable partial and total orders. *Z. Math. Logik Grundlagen Math.*, 22(1):1–18, 1976.
- [3] Rodney G. Downey, Denis R. Hirschfeldt, Steffen Lempp, and Reed Solomon. Computability-theoretic and proof-theoretic aspects of partial and linear orderings. *Israel J. Math.*, 138:271–289, 2003.
- [4] Noam Greenberg, Antonio Montalbán, and Theodore A. Slaman. Relative to any non-hyperarithmetical set. 2011. Submitted.
- [5] Denis R. Hirschfeldt, Bakhadyr Khoussainov, Richard A. Shore, and Arkadii M. Slinko. Degree spectra and computable dimensions in algebraic structures. *Ann. Pure Appl. Logic*, 115(1-3):71–113, 2002.
- [6] Denis R. Hirschfeldt and Richard A. Shore. Combinatorial principles weaker than Ramsey’s theorem for pairs. *J. Symbolic Logic*, 72(1):171–206, 2007.
- [7] Carl G. Jockusch, Jr., Bart Kastermans, Steffen Lempp, Manuel Lerman, and Reed Solomon. Stability and posets. *J. Symbolic Logic*, 74(2):693–711, 2009.
- [8] Julia F. Knight. Degrees coded in jumps of orderings. *J. Symbolic Logic*, 51(4):1034–1042, 1986.

- [9] Dev K. Roy. Recursive versus recursively enumerable binary relations. *Studia Logica*, 52(4):587–593, 1993.
- [10] Stephen G. Simpson. *Subsystems of second order arithmetic*. Perspectives in Logic. Cambridge University Press, Cambridge, second edition, 2009.
- [11] Theodore A. Slaman. Relative to any nonrecursive set. *Proc. Amer. Math. Soc.*, 126(7):2117–2122, 1998.
- [12] Robert I. Soare. *Recursively enumerable sets and degrees*. Perspectives in Mathematical Logic. Springer-Verlag, Berlin, 1987. A study of computable functions and computably generated sets.
- [13] Stephan Wehner. Enumerations, countable structures and Turing degrees. *Proc. Amer. Math. Soc.*, 126(7):2131–2139, 1998.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NOTRE DAME, NOTRE DAME, INDIANA 46556
U.S.A.

E-mail address: cholak@nd.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NOTRE DAME, NOTRE DAME, INDIANA 46556
U.S.A.

E-mail address: ddzhafar@nd.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, BERKELEY, CALIFORNIA 94720
U.S.A.,

E-mail address: schweber@math.berkeley.edu

DEPARTMENT OF MATHEMATICS, CORNELL UNIVERSITY, ITHACA, NY 14853 U.S.A.,

E-mail address: shore@math.cornell.edu